

Gold-catalyzed intermolecular alkyne oxyarylation for C3-functionalization of benzothiophenes

Rist, Paige; Grainger, Richard; Davies, Paul

DOI:

[10.1021/acs.orglett.0c03596](https://doi.org/10.1021/acs.orglett.0c03596)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Rist, P, Grainger, R & Davies, P 2021, 'Gold-catalyzed intermolecular alkyne oxyarylation for C3-functionalization of benzothiophenes', *Organic Letters*, vol. 23, no. 3, pp. 642–646.
<https://doi.org/10.1021/acs.orglett.0c03596>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Gold-Catalyzed Intermolecular Alkyne Oxyarylation for C3 Functionalization of Benzothiophenes

Paige A. Rist, Richard S. Grainger,* and Paul W. Davies*



Cite This: *Org. Lett.* 2021, 23, 642–646



Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information



ABSTRACT: C3-selective C–C bond formation on benzothiophenes is challenging, and few direct functionalization methods are available. A gold-catalyzed reaction of alkynes with benzothiophene S-oxides provides regioselective entry into C3-alkylated benzothiophenes with the C7-alkylated isomer as the minor product. This oxyarylation reaction works with alkyl and aryl alkynes and substituted and unsubstituted benzothiophenes. Mechanistic studies identify that sulfoxide inhibits the catalyst [DTBPAu(PhCN)]SbF₆, which also degrades and forms the unreactive complex [(DTBP)₂Au]SbF₆.

Benzothiophene-containing molecules display unique photophysical and electrochemical properties¹ as well as an array of biological activities against diseases such as cancer, HIV, diabetes, and Alzheimer's disease, rendering them of interest to the materials and pharmaceutical sectors (Figure 1).²

The direct functionalization of a benzothiophene provides a synthetically appealing entry into more substituted species, yet diverse methods for selective and direct C–C bond formation at C3 are scarce.³ Examples include palladium-catalyzed C–H arylations, which require relatively forcing conditions (Scheme 1a),⁴ and single examples of alkynylations⁵ and vinylations.⁶





Figure 1. Biologically important molecules that contain C3-substituted benzothiophenes.


Scheme 1. Selective Benzothiophene C3 Elaboration

a) Current methods


Palladium-catalyzed arylation - ref. 4



Remote Pummerer-type arylation and allylation - ref. 7




b) This work Gold-catalyzed alkylation



Received: October 28, 2020

Published: January 19, 2021



Procter's group introduced an interrupted Pummerer approach allowing C3 arylation and allylation processes on benzothiophene S-oxides (Scheme 1a).⁷ Here we report a mechanistically distinct benzothiophene functionalization method that uses simple alkynes to access complementary functionality at C3 through a gold-catalyzed oxyarylation process (Scheme 1b).

Gold- and subsequently Brønsted acid-catalyzed C–C bond-forming rearrangements between sulfoxides and alkynes have shown great promise for atom-efficient aryl C–H functionalization and other transformations.⁸ For the desired transformation, the vinylgold carbenoid intermediate **B** should evolve through a [3,3]-sigmatropic rearrangement and onto C3 rather than the C7 benzenoid position (Scheme 1b). Rearrangement must occur before the electrophilic organogold species reacts with a second equivalent of sulfoxide.^{8c} Our interest in gold-catalyzed reactions with sulfoxides⁹ led us to investigate the potential of oxyarylation for C3 benzothiophene elaboration.

Our study started with 2-methylbenzothiophene S-oxide (**1**) and 1-hexyne (**2a**), as unsubstituted benzothiophene S-oxide is unstable and decomposes out of solution (vide infra).¹⁰ All reactions were performed under non-inert conditions with undried, commercial solvents. A range of Au(I) catalysts was tested (Table 1, entries 1–6). In all cases the product **3a** from reaction at C3 was preferred, but some of the C7-functionalized product **4a** was also seen. Low reactivity occurred with SPhosAuNTf₂ and was not improved with higher temperatures

(Table 1, entries 1 and 2). Cleaner reactions and greater conversion to the desired product were seen with the phosphite-containing catalyst [DTBPAu(PhCN)]SbF₆ over those derived from more electron-rich ligands, with small amounts of benzothiophene **5** also observed in the latter case (Table 1, entries 1–5). Use of a more coordinating tosylate counterion saw reduced activity (Table 1, entry 6).

Improved yields were obtained in CH₂Cl₂ and fluorobenzene (Table 1, entries 7 and 8). A more substantial solvent effect was seen upon switching from hexyne to phenylacetylene (**2b**), with fluorobenzene giving a superior outcome with a 90% combined yield of oxyarylation products and much higher regioselectivity (Table 1, entry 10 vs 11). A modest reduction in yield was seen at a catalyst loading of 2.5 mol % (Table 1, entry 12). Reducing the equivalents of alkyne lowered the overall yield of **3** and **4** with both 1-hexyne and phenylacetylene (Table 1, entries 7 vs 9 and 11 vs 13). Reversing the stoichiometry of sulfoxide **1** and phenylacetylene **2b** further reduced the combined yield of **3b** and **4b** (Table 1, entry 14).


The best conditions were then applied across a range of alkynes and benzothiophene S-oxides (Scheme 2), with the reactions typically performed on a 0.2–0.5 mmol scale. On the 1.0 mmol scale, isomerically pure **3b** was isolated in 68% yield. Both electron-rich and electron-deficient aryl alkynes underwent oxyarylation successfully. Appreciable amounts of the free benzothiophene were observed with more electron-rich aryl alkynes (Scheme 2, compounds **6–9** and **24**). A low yield and poor C3:C7 regioselectivity were seen with an *o*-methoxy group on the aromatic ring. Aliphatic alkynes provided excellent yields with lower regioselectivities compared with aromatic alkynes (Scheme 2, compounds **15–18** and **23**). Good functional group tolerance was shown across these reactions with aryl halides, primary alkyl bromide, a tertiary amine, phthalimide, and carboxylic esters all readily incorporated.

The formation of **12–15** and **25** (Scheme 2) required higher temperatures for oxyarylation to occur. During the formation of **15** with 2.0 equiv of alkyne, the alkyne hydration product was also formed and was inseparable from the desired product. Using 1.0 equiv of alkyne saw no formation of the hydration side product, and **15** was isolated in good yield. No oxyarylation product was formed when diphenylacetylene and **1** were submitted to the reaction conditions. Only the benzil product from double oxidation of the alkyne was isolated (98% based on sulfoxide **1**).

Benzothiophene S-oxides with different substitution patterns also performed well. The reaction proceeds well in the absence of a C2 substituent, and halogen substituents are also tolerated (Scheme 2, **19–25**). In several cases very high regioselectivities were observed (>20:1), so simple trituration with methanol was sufficient to obtain the single C3 regioisomer (Scheme 2, **21**, **24**, and **25**). Good outcomes were also observed with an ester group at C2 (Scheme 2, **26** and **27**). The presence of a substituent at C3 did not lead to efficient formation of the C7 oxyarylation product, as a complex mixture was formed upon reaction of 3-methylbenzothiophene S-oxide with phenylacetylene **2b**.


As unsubstituted benzothiophene S-oxide is unstable when neat, a protocol was developed that telescoped together the S-oxidation of benzothiophene with the gold-catalyzed oxyarylation. C3-substituted benzothiophene **28** was obtained in 79% yield over two steps (see the Supporting Information for details).

Table 1. Reaction Optimization Study between 2-Methylbenzothiophene S-Oxide and Hex-1-yne or Phenylacetylene



entry	2 (equiv)	catalyst ^a	solvent	yields (1 : 3 : 4 : 5) (%) ^b
1	2a (2)	A	PhMe	70:16:2:0
2	2a (2)	A (23 °C)	PhMe	74:16:2:0
3	2a (2)	B	PhMe	32:39:10:— ^c
4	2a (2)	C	PhMe	46:33:4:— ^c
5	2a (2)	D	PhMe	— ^d :55:23:0
6	2a (2)	E /AgOTs ^e	PhMe	88:6:4:0
7	2a (2)	D	CH ₂ Cl ₂	0:64:23:— ^c
8	2a (2)	D	C ₆ H ₅ F	7:63:24:0
9	2a (1)	D	CH ₂ Cl ₂	0:50:17:<6 ^f
10	2b (2)	D	CH ₂ Cl ₂	— ^d :51:9:— ^c
11	2b (2)	D	C ₆ H ₅ F	0:80:10:<7 ^f
12	2b (2) ^g	D (2.5 mol %)	C ₆ H ₅ F	0:67:12:<10 ^f
13	2b (1)	D	C ₆ H ₅ F	0:62:9:8
14	2b (1) ^g	D	C ₆ H ₅ F	— ^d :50:9:— ^c

^a**A**, SPhosAuNTf₂; **B**, IPrAuNTf₂; **C**, [JohnPhosAu(MeCN)]SbF₆; **D**, [DTBPAu(PhCN)]SbF₆; **E**, DTBPAuCl; DTBP = (2,4-(^tBu)₂C₆H₃O)₃P. ^bYields and ratios were determined by ¹H NMR spectroscopy using a known concentration of 1,2,4,5-tetramethylbenzene. ^c**5** was present, but the yield could not be determined because of overlapping resonances. ^d**1** was present, but the yield could not be determined because of overlapping resonances. ^e**E** (0.05 equiv) and AgOTs (0.10 equiv). ^fRepresents the maximum possible yield because of overlapping resonances. ^g2 equiv of **1**.

Scheme 2. Scope of Gold-Catalyzed Oxyarylation of Alkynes with Benzothiophene S-Oxides^f

^aThe reaction was performed on a 1.0 mmol scale and afforded the C3:C7 mixture in 84% yield; the pure C3 isomer **3b** was isolated by column chromatography and recrystallization in 68% yield. ^bThe reaction was run at 75 °C. ^cThe reaction was run at 70 °C. ^dThe reaction was run at 50 °C. ^eThe reaction was run at 50 °C with 1.0 equiv of alkyne. ^fRatios refer to the isolated mixtures of C3:C7 regioisomers. Where ratios are not given, the yield refers to the isolated yield of the C3 regioisomer. The standard conditions used, unless stated otherwise, are 2.0 equiv of alkyne, 5 mol % [DTBPAu(PhCN)]SbF₆, and a reaction temperature of 0 °C.

Studies were then initiated in order to gain further insight into the mechanism (Scheme 3 and further details in the Supporting Information). Crossover experiments saw the formation of the oxyarylation products from only the benzothiophene S-oxide component and not the benzothiophene dopant (Scheme 3a).

Higher effective equivalents of sulfoxide **1** gave reduced yields of the functionalized benzothiophenes (Table 1, entries 9, 13, and 14). Monitoring the progress of an oxyarylation reaction of **1** and **2a** by in situ ¹H NMR spectroscopy showed the significant effect of the sulfoxide concentration (Scheme 3b). Under otherwise identical conditions, doubling the concentration of the limiting sulfoxide reagent **1** led to a substantially lower concentration of products **3b/4b** over the same reaction time (Scheme 3b), with no increase in the formation of free benzothiophene **5**. No other side products were seen in either reaction. A simultaneous comparison of batch- and slow-addition of the sulfoxide showed that the latter yielded more oxyarylation products **3b/4b** (Scheme 3b). An increase in the yield of oxyarylation products with more equivalents of alkyne was also observed (Table 1, entries 7 vs 9 and 11 vs 13). No reduction in reaction progress was seen



when a catalysis reaction was doped with an oxyarylation product (Scheme 3c).

An upfield shift in the ³¹P NMR resonance from 88 to 81 ppm was observed upon addition of sulfoxide **1** to [DTBPAu(PhCN)]SbF₆ at 0 °C. The free DTBP ligand was not observed in the ³¹P NMR or ¹H NMR spectra, while benzonitrile was observed in the ¹H NMR spectrum. A new minor resonance at 120 ppm was also observed, but this could also be seen in small amounts from the untreated catalyst in CDCl₃. Addition of phenylacetylene to [DTBPAu(PhCN)]SbF₆ resulted in the formation of several new ³¹P resonances, with a major one at 120 ppm. This resonance was identified as [(DTBP)₂Au]SbF₆ by in situ synthesis and also by isolation (Scheme 3d, i). The bisphosphite complex was formed preferentially and irreversibly in the presence of free ligand, with no change observed on addition of excess benzonitrile, sulfoxide, or phenylacetylene (Scheme 3d, ii). Using [(DTBP)₂Au]SbF₆ in a reaction with sulfoxide **1** and phenylacetylene gave only a trace amount of product (<5%) after an extended reaction time.



On the basis of these observations, the broader mechanism is consistent with that established by Ujaque et al.,^{8b} where sulfoxide addition to an alkyne–gold complex generates a

Scheme 3. Mechanistic Studies

a) Crossover experiments

b) *In situ* ^1H NMR spectroscopic studies into the effect of sulfoxide concentration

c) Product-doped inhibition experiment

d) ^{31}P NMR spectroscopic studies into catalyst deactivation

vinylgold carbenoid intermediate that undergoes [3,3]-sigmatropic rearrangement and then rearomatization and protodeauration. The alternative carbene formation and electrophilic aromatic substitution pathway can be ruled out from the results of the crossover experiments and the ready formation of compounds 26 and 27 bearing a deactivating ester group at C2.

Sulfoxide inhibition of the reaction is seen, as are direct interactions between sulfoxide 1 and the gold complex. The release of DTBP by catalyst degradation pathways, as observed in the presence of phenylacetylene, will reduce the amount of any remaining active catalyst by formation of $[(\text{DTBP})_2\text{Au}]\text{SbF}_6$, which is essentially a catalytic dead-end for the desired process.

In summary, gold-catalyzed reactions between readily accessed benzothiophene S-oxides and terminal alkynes provide efficient and selective access to C3-alkylated benzothiophenes bearing a useful carbonyl group for subsequent exploitation. The approach is compatible with a variety of electron-rich and electron-deficient aromatic and aliphatic alkynes as well as reactive functionalities including carboxylic esters, tertiary amines, phthalimide, and alkyl and aryl halides. The highest regioselectivities are seen with aryl alkynes. The reactions are straightforward to perform and do

not require dry or inert conditions or high temperatures. This is the first instance of the use of gold-catalyzed oxyarylation to modify a sulfur-containing ring. The reaction is compatible with substitution around the benzothiophene, including halogens and electron-withdrawing groups. A telescoped oxidation–oxyarylation sequence has also been developed.

Sulfoxide was determined to inhibit the gold-catalyzed process, which may have wider relevance in other gold-catalyzed transformations. This study also highlights the need for further development of more robust gold complexes that retain the reactivity profile of phosphite Au(I) species.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.0c03596>.

Details of the effect of the sulfoxide 1 concentration on oxyarylation reactions, *in situ* ^{31}P NMR catalyst studies, assignment of regioisomers 3b and 4b, experimental details, compound characterization data, and copies of NMR spectra (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

Paul W. Davies – School of Chemistry, University of Birmingham, Birmingham B15 2TT, U.K.; orcid.org/0000-0002-0340-2414; Email: p.w.davies@bham.ac.uk

Richard S. Grainger – School of Chemistry, University of Birmingham, Birmingham B15 2TT, U.K.; orcid.org/0000-0003-0611-4386; Email: r.s.grainger@bham.ac.uk

Author

Paige A. Rist – School of Chemistry, University of Birmingham, Birmingham B15 2TT, U.K.

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.orglett.0c03596>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the EPSRC for a studentship to P.A.R. (EP/N509590/1). The authors gratefully acknowledge support from the Centre for Chemical and Materials Analysis in the School of Chemistry and Dr. Cécile le Duff for NMR support. P.W.D. is grateful to the Royal Society and the Leverhulme Trust for the award of a Senior Research Fellowship (SRF\R1\191033).

■ REFERENCES

- (a) Takimiya, K.; Shinamura, S.; Osaka, I.; Miyazaki, E. Thienocene-Based Organic Semiconductors. *Adv. Mater.* **2011**, *23*, 4347–4370. (b) Sumi, T.; Takagi, Y.; Yagi, A.; Morimoto, M.; Irie, M. Photoirradiation wavelength dependence of cycloreversion quantum yields of diarylethenes. *Chem. Commun.* **2014**, *50*, 3928–3930. (c) Ruzié, C.; Karpinska, J.; Laurent, A.; Sanguinet, L.; Hunter, S.; Anthopoulos, T. D.; Lemaire, V.; Cornil, J.; Kennedy, A. R.; Fenwick, O.; Samorì, P.; Schweicher, G.; Chattopadhyay, B.; Geerts, Y. H. Design, synthesis, chemical stability, packing, cyclic voltammetry, ionisation potential, and charge transport of [1]benzothieno[3,2-*b*]-[1]benzothiophene derivatives. *J. Mater. Chem. C* **2016**, *4*, 4863–4879. (d) Benincori, T.; Gámez-Valenzuela, S.; Goll, M.; Bruchlos, K.

- Malacrida, C.; Arnaboldi, S.; Mussini, P. R.; Panigati, M.; López Navarrete, J. T.; Ruiz Delgado, M. C.; Appoloni, G.; Ludwigs, S. Electrochemical studies of a new, low-band gap inherently chiral ethylenedioxythiophene-based oligothiophene. *Electrochim. Acta* **2018**, *284*, 513–525. (e) Kim, D.; Kwon, J. E.; Park, S. Y. Fully Reversible Multistate Fluorescence Switching: Organogel System Consisting of Luminescent Cyanostilbene and Turn-On Diarylethene. *Adv. Funct. Mater.* **2018**, *28*, 1706213–1706219. (f) Usta, H.; Kim, D.; Ozdemir, R.; Zorlu, Y.; Kim, S.; Ruiz Delgado, M. C.; Harbuzaru, A.; Kim, S.; Demirel, G.; Hong, J.; Ha, Y.-G.; Cho, K.; Facchetti, A.; Kim, M. High Electron Mobility in [1]Benzothieno[3,2-*b*][1]-benzothiophene-Based Field-Effect Transistors: Toward *n*-Type BTBTs. *Chem. Mater.* **2019**, *31*, 5254–5263.
- (2) (a) Chapman, N. B.; Scrowston, R. M.; Westwood, R. Pharmacologically Active Benzo[*b*]thiophen Derivatives. Part VIII. Benzo[*b*]thiophen Analogues of Tryptophan and α -Methyltryptophan, and Some of their 5-Substituted Derivatives. *J. Chem. Soc. C* **1969**, *0*, 1855–1857. (b) Muchmore, D. B. Raloxifene: A Selective Estrogen Receptor Modulator (SERM) with Multiple Target System Effects. *Oncologist* **2000**, *5*, 388–392. (c) Willoughby, C. A.; Bull, H. G.; García-Calvo, M.; Jiang, J.; Chapman, K. T.; Thornberry, N. A. Discovery of Potent, Selective Human Granzyme B Inhibitors that Inhibit CTL Mediated Apoptosis. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 2197–2200. (d) Croxtall, J. D.; Plosker, G. L. Sertaconazole. *Drugs* **2009**, *69*, 339–359. (e) Boitano, A. E.; Wang, J.; Romeo, R.; Bouchez, L. C.; Parker, A. E.; Sutton, S. E.; Walker, J. R.; Flaveny, C. A.; Perdew, G. H.; Denison, M. S.; Schultz, P. G.; Cooke, M. P. Aryl Hydrocarbon Receptor Antagonists Promote the Expansion of Human Hematopoietic Stem Cells. *Science* **2010**, *329*, 1345–1348. (f) Chiummiento, L.; Funicello, M.; Lupattelli, P.; Tramutola, F.; Berti, F.; Marino-Merlo, F. Synthesis and biological evaluation of novel small non-peptidic HIV-1 PIs: The benzothiophene ring as an effective moiety. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 2948–2950. (g) Hansen, A. H.; Sergeev, E.; Pandey, S. K.; Hudson, B. D.; Christiansen, E.; Milligan, G.; Ulven, T. Development and Characterization of a Fluorescent Tracer for the Free Fatty Acid Receptor 2 (FFA2/GPR43). *J. Med. Chem.* **2017**, *60*, 5638–5645. (h) Keri, R. S.; Chand, K.; Budagumpi, S.; Balappa Somappa, S.; Patil, S. A.; Nagaraja, B. M. An overview of benzo[*b*]thiophene-based medicinal chemistry. *Eur. J. Med. Chem.* **2017**, *138*, 1002–1033.
- (3) Bheeter, C. B.; Chen, L.; Soulé, J.; Doucet, H. Regioselectivity in palladium-catalysed direct arylation of 5-membered ring heteroaromatics. *Catal. Sci. Technol.* **2016**, *6*, 2005–2049.
- (4) (a) Kirchberg, S.; Tani, S.; Ueda, K.; Yamaguchi, J.; Studer, A.; Itami, K. Oxidative biaryl coupling of thiophenes and thiazoles with arylboronic acids through palladium catalysis: otherwise difficult C4-selective C-H arylation enabled by boronic acids. *Angew. Chem., Int. Ed.* **2011**, *50*, 2387–2391. (b) Biajoli, A. F. P.; da Penha, E. T.; Correia, C. R. D. Palladium catalysed regioselective arylation of indoles, benzofuran and benzothiophene with aryl diazonium salts. *RSC Adv.* **2012**, *2*, 11930–11935. (c) Tang, D. D.; Collins, K. D.; Ernst, J. B.; Glorius, F. Pd/C as a Catalyst for Completely Regioselective C-H Functionalization of Thiophenes under Mild Conditions. *Angew. Chem., Int. Ed.* **2014**, *53*, 1809–1813. (d) Gong, W.-J.; Liu, D.-X.; Li, F.-L.; Gao, J.; Li, H.-X.; Lang, J.-P. Palladium-catalyzed decarboxylative C3-acylation of benzofurans and benzothiophenes with α -oxocarboxylic acids via direct sp^2 C-H bond activation. *Tetrahedron* **2015**, *71*, 1269–1275. (e) Colletto, C.; Islam, S.; Juliá-Hernández, F.; Larrosa, I. Room-Temperature Direct β -Arylation of Thiophenes and Benzo[*b*]thiophenes and Kinetic Evidence for a Heck-type Pathway. *J. Am. Chem. Soc.* **2016**, *138*, 1677–1683. (f) Tang, D. D.; Collins, K. D.; Glorius, F. Completely regioselective direct C-H functionalization of benzo[*b*]thiophenes using a simple heterogeneous catalyst. *J. Am. Chem. Soc.* **2013**, *135*, 7450–7453.
- (5) Mahato, S. K.; Chatani, N. The Iridium(III)-Catalyzed Direct C(sp^2)- and C(sp^3)-H Alkynylation of 2-Acylimidazoles with Various Alkynyl Bromides: Understanding the Full Catalytic Cycle. *ACS Catal.* **2020**, *10*, 5173–5178.
- (6) (a) Ueyama, T.; Mochida, S.; Fukutani, T.; Hirano, K.; Satoh, T.; Miura, M. Ruthenium-Catalyzed Oxidative Vinylation of Heteroarene Carboxylic Acids with Alkenes via Regioselective C-H Bond Cleavage. *Org. Lett.* **2011**, *13*, 706–708. (b) Zhang, Y.; Li, Z.; Liu, Z. Pd-Catalyzed Olefination of Furans and Thiophenes with Allyl Esters. *Org. Lett.* **2012**, *14*, 226–229.
- (7) Shriver, H. J.; Fernández-Salas, J. A.; Hedtke, C.; Pulis, A. P.; Procter, D. J. Regioselective synthesis of C3 alkylated and arylated benzothiophenes. *Nat. Commun.* **2017**, *8*, 14801.
- (8) (a) Shapiro, N. D.; Toste, F. D. Rearrangement of Alkynyl Sulfoxides Catalyzed by Gold(I) Complexes. *J. Am. Chem. Soc.* **2007**, *129*, 4160–4161. (b) Cuenca, A. B.; Montserrat, S.; Hossain, K. M.; Mancha, G.; Lledós, A.; Medio-Simón, M.; Ujaque, G.; Asensio, G. Gold(I)-Catalyzed Intermolecular Oxyarylation of Alkynes: Unexpected Regiochemistry in the Alkylation of Arenes. *Org. Lett.* **2009**, *11*, 4906–4909. (c) Lu, B.; Li, Y.; Wang, Y.; Aue, D. H.; Luo, Y.; Zhang, L. [3,3]-Sigmatropic Rearrangement versus Carbene Formation in Gold-Catalyzed Transformations of Alkynyl Aryl Sulfoxides: Mechanistic Studies and Expanded Reaction Scope. *J. Am. Chem. Soc.* **2013**, *135*, 8512–8524. (d) Barrett, M. J.; Davies, P. W.; Grainger, R. S. Regioselective functionalisation of dibenzothiophenes through gold-catalysed intermolecular alkyne oxyarylation. *Org. Biomol. Chem.* **2015**, *13*, 8676–8686. (e) Kaiser, D.; Veiros, L. F.; Maulide, N. Brønsted Acid-Mediated Hydrative Arylation of Unactivated Alkynes. *Chem. - Eur. J.* **2016**, *22*, 4727–4732. (f) Hu, L.; Gui, Q.; Chen, X.; Tan, Z.; Zhu, G. HOTf-Catalyzed, Solvent-Free Oxyarylation of Ynol Ethers and Thioethers. *J. Org. Chem.* **2016**, *81*, 4861–4868. (g) Baldassari, L. L.; Mantovani, A. C.; Senoner, S.; Maryasin, B.; Maulide, N.; Lüdtke, D. S. Redox-Neutral Synthesis of Selenoesters by Oxyarylation of Selenoalkynes under Mild Conditions. *Org. Lett.* **2018**, *20*, 5881–5885. (h) Li, G.; Zhang, L. Gold-Catalyzed Intramolecular Redox Reaction of Sulfinyl Alkynes: Efficient Generation of α -Oxo Gold Carbenoids and Application in Insertion into R-CO Bonds. *Angew. Chem., Int. Ed.* **2007**, *46*, 5156–5159. (i) Kaldre, D.; Maryasin, B.; Kaiser, D.; Gajsek, O.; González, L.; Maulide, N. An Asymmetric Redox Arylation: Chirality Transfer from Sulfur to Carbon through a Sulfonium [3,3]-Sigmatropic Rearrangement. *Angew. Chem., Int. Ed.* **2017**, *56*, 2212–2215.
- (9) (a) Barrett, M. J.; Khan, G. F.; Davies, P. W.; Grainger, R. S. Alkynyl sulfoxides as α -sulfinyl carbene equivalents: gold-catalysed oxidative cyclopropanation. *Chem. Commun.* **2017**, *53*, 5733–5736. (b) Davies, P. W.; Albrecht, S. J. C. Gold- or platinum-catalyzed synthesis of sulfur heterocycles: access to sulfur ylides without using sacrificial functionality. *Angew. Chem., Int. Ed.* **2009**, *48*, 8372–8375.
- (10) Pouzet, P.; Erdelmeier, I.; Ginderow, D.; Mornon, J.-P.; Dansette, P.; Mansuy, D. Thiophene S-Oxides: Convenient Preparation, First Complete Structural Characterization and Unexpected Dimerization of One of Them, 2,5-Diphenylthiophene-I-oxide. *J. Chem. Soc., Chem. Commun.* **1995**, 473–474.